

B<sup>1</sup> cont  
U.S. Application No. 09/071,838, filed May 1, 1998, each of which is incorporated by reference in its entirety.--

---

[Please amend the first paragraph on page 7 of the application as follows:]

B<sub>2</sub>  
Another example of algorithm that is suitable for determining percent sequence identity and sequence similarity is the BLAST algorithm, which is described in Altschul *et al.*, *J. Mol. Biol.* 215:403-410 (1990). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information. This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul *et al.*, *supra*). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLAST program uses as defaults a wordlength (W) of 11, the BLOSUM62 scoring matrix (*see* Henikoff & Henikoff, *Proc. Natl. Acad. Sci. USA* 89:10915 (1989)) alignments (B) of 50, expectation (E) of 10, M=5, N=-4, and a comparison of both strands.

---

IN THE CLAIMS:

Please cancel claims 4, 11, and 14.

Please amend claims 1, 2, 9, and 15 and add claims 25-28<sup>9</sup> as follows:

---

B<sub>3</sub>  
1. (Twice amended) An isolated double-stranded nucleic acid molecule comprising a *FIE* polynucleotide encoding a polypeptide at least 60% identical